

2018-2019年度河南省种子协会小麦品种比较试验汇总结果（旱地组）

| 组别 | 序号 | 品种名称 | 比CK晚熟(天) | 亩产(kg) | 比CK增减(%) | 增产点数 | 增产率(%) | 严重倒伏点率(%) | 总位次 | 条锈病 | 叶锈病 | 白粉病 | 纹枯病 | 赤霉病 |
|-----|----|--------|----------|--------|----------|------|--------|-----------|-----|-----|-----|-----|-----|-----|
| 旱地组 | 1 | 豫丰硕002 | 1.2 | 378.8 | 7.9 | 10 | 100.0 | 0.0 | 1 | MS | HR | MS | MS | HS |
| 旱地组 | 2 | 偃亳505 | 1.3 | 379.0 | 7.3 | 10 | 100.0 | 0.0 | 2 | MS | HR | MS | MR | MS |
| 旱地组 | 3 | 洛旱26 | -0.4 | 353.0 | 4.5 | 8 | 80.0 | 0.0 | 3 | MR | MR | MS | MS | HS |
| 旱地组 | 4 | 豫农523 | -0.1 | 361.2 | 4.5 | 9 | 90.0 | 0.0 | 3 | MS | MR | MS | MS | HS |
| 旱地组 | 5 | 洛旱33 | -0.6 | 352.0 | 4.2 | 8 | 80.0 | 0.0 | 5 | MS | MR | MS | MS | HS |
| 旱地组 | 6 | 豫迪999 | 0.0 | 365.0 | 4.0 | 8 | 80.0 | 0.0 | 6 | MS | HR | MR | MS | MS |
| 旱地组 | 7 | 中育1215 | 0.5 | 363.4 | 2.9 | 6 | 60.0 | 0.0 | 7 | MR | HR | MS | MS | HS |
| 旱地组 | 8 | 中麦41 | 1.4 | 345.8 | 2.3 | 7 | 70.0 | 0.0 | 8 | HS | HR | MS | MS | HS |
| 旱地组 | 9 | 宛麦168 | 1.2 | 352.3 | 1.1 | 6 | 60.0 | 0.0 | 9 | MR | MS | MS | MS | MS |
| 旱地组 | 10 | 偃高163 | 1.0 | 347.0 | 0.3 | 6 | 60.0 | 0.0 | 10 | MS | MS | MS | MS | HS |
| 旱地组 | 11 | 金麦996 | 1.1 | 349.7 | -0.4 | 6 | 60.0 | 0.0 | 11 | MS | MR | MR | MS | HS |
| 旱地组 | 12 | 焦麦278 | 0.5 | 343.9 | -2.0 | 4 | 40.0 | 0.0 | 12 | MR | MR | MR | MS | HS |
| 旱地组 | 13 | 佳源339 | 1.3 | 343.3 | -2.2 | 5 | 50.0 | 0.0 | 13 | MS | HR | MS | MS | HS |
| 旱地组 | 14 | 智优343 | 0.9 | 338.2 | -2.2 | 4 | 40.0 | 0.0 | 13 | MR | HR | MS | HR | HS |
| 旱地组 | 15 | 郑丰136 | 1.6 | 337.9 | -2.3 | 5 | 50.0 | 0.0 | 15 | MR | HR | MS | MR | HS |
| 旱地组 | 16 | 偃麦138 | 0.9 | 339.4 | -3.9 | 3 | 30.0 | 0.0 | 16 | MS | HR | MS | MR | HS |
| 旱地组 | 17 | 河大357 | 0.3 | 324.5 | -4.0 | 4 | 40.0 | 0.0 | 17 | MS | HR | MS | HR | HS |
| 旱地组 | 18 | 宇宝881 | 0.3 | 336.3 | -4.2 | 1 | 10.0 | 0.0 | 18 | MR | HR | MS | MR | HS |
| 旱地组 | 19 | 才智911 | 1.3 | 328.8 | -4.9 | 6 | 60.0 | 0.0 | 19 | MS | MR | MS | MS | HS |
| 旱地组 | 20 | 仁麦20 | 0.1 | 328.1 | -5.1 | 3 | 30.0 | 0.0 | 20 | MR | MR | MS | MS | HS |
| 旱地组 | 21 | 洛新183 | 1.5 | 332.6 | -5.9 | 3 | 30.0 | 0.0 | 21 | MS | MS | MR | MS | HS |
| 旱地组 | 22 | 天民319 | -0.1 | 329.8 | -6.6 | 2 | 20.0 | 0.0 | 22 | MR | HR | HR | MS | HS |
| 旱地组 | 23 | 中科124 | 1.6 | 314.5 | -6.9 | 2 | 20.0 | 0.0 | 23 | MS | HR | MS | MS | HS |
| 旱地组 | 24 | 田丰182 | -0.6 | 327.2 | -7.4 | 2 | 20.0 | 0.0 | 24 | MS | MS | MR | MS | HS |
| 旱地组 | 25 | 广延257 | -0.1 | 311.2 | -7.9 | 1 | 10.0 | 0.0 | 25 | MR | MR | MR | MS | HS |
| 旱地组 | 26 | 牟农181 | 1.7 | 310.3 | -8.2 | 5 | 50.0 | 0.0 | 26 | HS | HR | MS | HR | MS |
| 旱地组 | 27 | 鹤麦1704 | 0.6 | 309.9 | -8.3 | 2 | 20.0 | 0.0 | 27 | MS | HR | MS | MS | HS |
| 旱地组 | 28 | 偃科508 | 1.2 | 312.4 | -9.7 | 4 | 40.0 | 0.0 | 28 | HS | HS | MS | MS | HS |
| 旱地组 | 29 | 安工5号 | 1.1 | 316.8 | -9.7 | 2 | 20.0 | 0.0 | 28 | HS | MS | MR | MS | HS |
| 旱地组 | 30 | 郑麦180 | 0.5 | 308.9 | -10.7 | 1 | 10.0 | 0.0 | 30 | MS | HR | MS | MS | MS |
| 旱地组 | 31 | 盛贤1号 | -0.1 | 308.7 | -12.0 | 0 | 0.0 | 0.0 | 31 | MS | HR | MR | MS | MS |

抗病鉴定结果： HR:高抗； MR:中抗； HS:高感； MS:中感
 严重倒伏点率：倒伏程度 ≥ 4 级且倒伏面积 $\geq 30.0\%$ 的试点率
 比CK增减数值保留1位小数后相同的品种，总位次并列。